

MISDIRECTED AMPLEXUS BETWEEN A PACIFIC TREEFROG (*PSEUDACRIS REGILLA*) AND A WESTERN TOAD (*ANAXYRUS BOREAS*) IN A NORTHERN CALIFORNIA UPLAND

JEFF A ALVAREZ, ELIZABETH LOPEZ, TARA COLLINS, ANGELIQUE HERMAN, JENNIFER JELINCIC, MARINA L OLSON, BRIAN PIONTEK, ALICIA I RINGSTAD, AND JEFFERY T WILCOX

ABSTRACT—Misdirected amplexus has been reported between conspecifics, congeners, between different amphibian families, and between amphibians and non-amphibian objects. We observed misdirected amplexus between a Pacific Treefrog and a Western Toad, wherein the treefrog amplexed the toad's leg. These types of events can result in reproductive interference and have also been considered an evolutionary trap for some species.

Key words: amplexus, evolutionary trap, misdirected, Pacific Treefrog, reproductive interference, Western Toad

Misdirected amplexus has been defined variously as amplexus between 2 individuals of similar species (for example, *Rana boylei* and *Rana catesbeiana*; Storm 1952; Lind and others 2003; Pearl and others 2005), 2 different genera (for example, *Bufo boreas* and *Rana draytonii*; Alvarez 2011; Machado and Bernarde 2011), 2 species of different families (for example, Caudata and Anura; Waterstrat and others 2008; Simovic' and others 2014), and amplexus between an anuran and inanimate objects (Storm 1960; Brown 1977). These types of amplexal behaviors are attributed to a strong drive for reproduction, often with a much larger available female (Schmeller and others 2005). Misdirected amplexus can also occur intraspecifically in that male-male amplexus has been reported by Bowcock and others (2008), as well as other researchers. In many anurans, release calls and other physical movements are used to reject unwanted or erroneous amplexus by unreceptive females or males, but these behavioral cues are likely only discouraging to conspecific males (Stebbins and Cohen 1995).

On 4 March 2019, while conducting a California Red-legged Frog (*Rana draytonii*) egg mass survey at the Mitsui Ranch (UTM: Zone 10S, 536751.9 m E, 4241885.4 m N; determined using Etrex handheld GPS unit, Garmin Limited, Lenexa, Kansas, USA) with numerous col-

leagues, a partially buried rock functioning as an upland cover object was lifted, and within a small cavity below was an adult Western Toad (*Anaxyrus boreas*). We removed the Western Toad for closer inspection and noted that a Pacific Treefrog (*Pseudacris regilla*) appeared to be clinging to the posterior portion of the toad. When inspected closely, it was clear that the Pacific Treefrog was in amplexus with the tibiofibula portion of the hind left leg (posterior portion) of the toad just above the tarsal bone (Fig. 1). The absence of nuptial pads on the toad helped us to identify it as a female, while the treefrog was clearly identified as a male due the skin folds of the relaxed vocal sacs. The treefrog remained amplexed with the Western Toad for several minutes while the toad was handled and photographed. Shortly before release, the treefrog loosened its grasp and remained in a resting position on the dorsum of the toad. Thereafter, both toad and treefrog were returned to the site of capture and the rock was replaced as found.

Misdirected amplexus events are not rare, but it is unusual in animals exhibiting such size dimorphism as in this observation. The difference in size between the 2 individuals almost certainly explains the location of the amplexus. Most amplexus is thoracic or axial, and species appear to utilize a specific type or place for amplexus (Duellman and Trueb 1994; Stebbins and Cohen 1995).

Generally, misdirected amplexus can create reproductive interference, which can reduce the fitness of individual animals at some locations (Pearl and others 2005; D'Amore and others 2009). Misdirected and intraspecific amplexus can also become an evolutionary trap for declining species (see D'Amore and others 2009), although in species that are not in decline the individuals may only lose a breeding season or, in some cases, increase their own exposure to predation (Alvarez 2011).



FIGURE 1. Adult female Western Toad (*Anaxyrus boreas*) amplexed by an adult male Pacific Treefrog (*Pseudacris regilla*), Sonoma County, California, 4 March 2019 (Photo by M Olsen).

We speculate that the likely result of an event such as the one described above is short-term sequestration of the individuals from reproductive activity. We can only assume that had we not interfered, the toad would have left the shelter of the rock and moved overland to a breeding site several meters away. We speculate that in the event the treefrog had been able to maintain amplexus during this short overland migration, once in the pond the treefrog may have been dislodged by 1 or more male toads attempting to amplex the female toad. However, this misdirected amplexus could also have resulted in increased predation potential for 1 or both individuals since natural movement could be impacted (Alvarez 2011). The true nature of the effects of these types of events has yet to be studied unambiguously, and their inconsistent frequency at most sites will likely contribute to difficulty in interpreting the effects of misdirected amplexus.

Acknowledgements.—We wish to thank the Mitsui Ranch and the Sonoma Mountain Ranch Preservation Foundation for access to the site, and for continued support to conduct surveys for amphibians. We also thank J Bowerman for editorial comments that improved the manuscript.

LITERATURE CITED

- ALVAREZ JA. 2011. *Bufo boreas* (Western Toad). Davian Behavior. *Herpetological Review* 42:408–409.
- BOWCOCK H, BROWN GP, SHINE R. 2008. Sexual communication in Cane Toads, *Chaunus marinus*: What cues influence the duration of amplexus? *Animal Behaviour* 75:1571–1579.
- BROWN H. 1977. A case of interbreeding between *Rana aurora* and *Bufo boreas* (Amphibia, Anura). *Journal of Herpetology* 11:92–94.
- D'AMORE A, KIRBY E, HEMINGWAY V. 2009. Reproductive interference by an invasive species: An evolutionary trap? *Herpetological Conservation and Biology* 4:325–330.
- DUELLMAN WE, TRUEB L. 1994. *Biology of amphibians*. Baltimore, MD: Johns Hopkins University Press. 670 p.
- LIND AJ, BETTASO JB, YARNELL SM. 2003. *Rana boylei* (Foothill Yellow-legged Frog) and *Rana catesbeiana* (Bullfrog). Reproductive behavior. *Herpetological Review* 34:234–235.
- MACHADO RA, BERNARDE PS. 2011. Multiple and heterospecific amplexi between the toads *Rhaebo guttatus* and *Rhinella marina* (Anura: Bufonidae). *Herpetology Notes* 4:167–169.
- PEARL CA, HAYES MP, HAYCOCK R, ENGLER JD, BOWERMAN J. 2005. Observations of interspecific amplexus between western North American ranid frogs and the introduced American Bullfrog (*Rana catesbeiana*) and an hypothesis concerning breeding interference. *American Midland Naturalist* 154:126–134.
- SCHMELLER DS, O'HARA R, KOKKO H. 2005. Male adaptive stupidity: Male mating pattern in hybrid-ogenetic frogs. *Evolutionary Ecology Research* 7:1039–1050.
- SIMOVIC'A, ANDERSON N, ANDELKOVIC' M, GVOZDENOVIC' S, ĐORDEVIC' S. 2014. Unusual amplexuses between anurans and caudates. *Herpetology Notes* 7:25–29.
- STEBBINS RC, COHEN NW. 1995. *A natural history of amphibians*. Princeton, NJ: Princeton University Press. 316 p.
- STORM RM. 1952. Interspecific behaviour in *Rana aurora* and *Rana catesbeiana*. *Herpetologica* 8:108.
- STORM RM. 1960. Notes on the breeding biology of the Red-legged Frog (*Rana aurora aurora*). *Herpetologica* 16:251–259.
- WATERSTRAT FT, MCINTYRE AP, HAYES MP, PHILLIPS KM, CURRY TR. 2008. *Ascaphus truei* (Coastal Tailed Frog). Atypical amplexus. *Herpetological Review* 39:458.

The Wildlife Project, PO Box 188888, Sacramento, CA 95818 USA (JAA); jeff@thewildlifeproject.com; Sequoia Ecological Consulting, 156 Diablo Road, Suite 320, Danville, CA 94526 USA (EL); Westervelt Ecological Services, 600 North Market Blvd, Suite 3, Sacramento, CA 95834 USA (TC); AECOM, 1194 Pacific Street, Suite 204, San Luis Obispo, CA 93401 USA (AH); Jacobs, 300 Frank H. Ogawa Plaza, Suite 600, Oakland, CA 94612 USA (JJ); 7029 Treasure Way, Sacramento, CA 95831 USA (MLO);

Horizon Water and Environment, 266 Grand Ranch Preservation Foundation, 3124 Sonoma Avenue, Suite 210, Oakland, CA 94610 USA (BP); Mountain Road, Petaluma, CA 94954 USA (JTW). Jacobzoon and Associates, Inc., 117 Clara Avenue, Ukiah, CA 95482 USA (AIR); Sonoma Mountian Submitted 27 August 2020, accepted 12 October 2020. Corresponding Editor: Jay Bowerman.