

Prolapsed hemipenis in an adult Baja California Coachwhip (*Masticophis fuliginosus*) in the Sierra San Pedro Mártir, Baja California, México

Jeff A. Alvarez, The Wildlife Project, PO Box 188888, Sacramento, CA, USA; jeff@thewildlifeproject.com

Carlos Alvarado-Laguna, Ascent Environmental, 455 Capitol Mall, Suite 300, Sacramento, CA, USA

Jorge H. Valdez-Villavicencio, Conservación de Fauna del Noroeste, A.C. Ensenada, Baja CA, C.P. 22800, México

Elizabeth K. Lopez, Sequoia Ecological Consulting, Inc. 2110 K St., Suite 11, Sacramento, CA, USA

Rafael A. Lara-Resendiz, Conservación de Fauna del Noroeste, A.C. Ensenada, Baja California, C.P. 22800, México

The Baja California Coachwhip (*Masticophis fuliginosus*) is a large (> 2 m), diurnal snake that is common throughout all of Baja California peninsula and is found in a variety of habitats ranging from desert and scrublands to riparian and agricultural areas (Stebbins 2003). The species is often observed foraging for lizards among open patches of vegetated habitat where it moves quickly along the ground and captures lizards and other prey by direct pursuit (Ortenburger 1925, Thomson et al. 2016). *Masticophis fuliginosus* is in relative decline, due to land use changes within its range (Thomson et al. 2016), and natural history investigations focused on this species are required to enable land managers and biologists the information needed to appropriately manage the species.

We regularly encounter *M. fuliginosus* and numerous other species of snakes when we enter appropriate habitat in northwestern Baja California. Herein we report an observation of a *M. fuliginosus* that exhibited a prolapsed hemipenis while otherwise completely unharmed and unmolested.

While participating in a herpetofauna natural history workshop, we observed a dark phase adult *M. fuliginosus* basking within open scrub/chaparral habitat at Rancho El Potrero, Sierra San Pedro Mártir (30.918586 N, 115.645567 W; elev. 900 m). Upon approach (RAL), the snake immediately attempted to flee but was hand-collected. At the time of capture, it was quickly noted that the right hemipenis was prolapsed (Fig 1). The snake was very carefully placed in a cotton bag, and then the bag was placed in a location where it would be undisturbed. After approximately 20 minutes the snake was removed from the bag for examination. At that time of the snake's removal from the bag, the hemipenis was still everted and extremely engorged, which can be a typical condition, but only during copulatory behavior (Kelly 2002). Podhade and Harne (2014) suggested that a prolapsed hemipenis would likely desiccate and become necrotic if not returned to its natural inverted position. One of the authors (JAA) very carefully attempted to slowly replace the hemipenis into its natural position, but it was clear that the hypertension

was too high for the hemipenis to be compressed and replaced in a manner that was safe for the snake.

Prolapsed hemipenes do occur through several reptilian groups. Vyas and Mistry (2021) noted that a dead, wild Marsh Crocodile (*Crocodylus palustris*) had a prolapsed phallus/cloaca in India. Lescano and Quevedo (2016) conducted a hemipenectomy on a captive Leopard Gecko (*Eublepharis macularius*), with successful survival of the lizard. A captive Central Asian Cobra (*Naja oxiana*) had a prolapsed hemipenis, which was subsequently amputated to avoid necrosis and infection (Nasoori et al. 2018). Prolapsed hemipenes are not unique to our observation; Bennett (1996) and Hernandez-Divers (2004) both reported on suggested surgical repairs for such conditions in several reptile groups (i.e., turtles, lizards, snakes).

It appears that wild specimens are seldom reported with a prolapsed hemipenis. We found that only 3 accounts worldwide, all recent: Vyas and Mistry (2021) reported this phenomenon in a dead wild crocodile (India), Pernetta and Reading (2021) collected a live snake in this condition (Australia), and Schultz and Caven (2021) reported it in a snake



Fig. 1. Prolapsed hemipene in an adult *Masticophis fuliginosus* in Baja California. Photo by Elizabeth Lopez.

The Baja California coachwhip (*Masticophis fuliginosus*) is a large (> 2 m), diurnal snake that is common throughout all of Baja California peninsula and is found in a variety of habitats ranging from desert and scrublands to riparian and agricultural areas (Stebbins 2003).

killed in a fire (Nebraska, USA). This paucity of reports may be due to a reduced ability to remain mobile, infection and necrosis to the prolapsed area, or other factors that reduce fitness of the specimen, perhaps leading to predation or death. Prolapse of the cloacal region, in both male and female reptiles, has been reported among many reptile groups (see Hedley and Eatwell 2014). However, very few reports of wild caught reptiles have been reported with a prolapsed hemipenis. Researchers have suggested reasons how a hemipenis could become prolapsed and included: parasitic loads, traumatic injury, chronic sexual activity, forced separation during copulation, trauma involving the retractor muscle of hemipenis or cloacal sphincter, regional infection within the individual animal, and other factors (De Silva et al. 2013, Podhade and Harne 2014, Schultz and Caven 2021). In the case of our observation, we speculate that the male sex organ may have been forcibly removed during copulation and was unable to be retracted. Due to the remoteness of our observation, any surgical/veterinary intervention was not practicable and therefore we released the snake at the site of capture with its hemipenis prolapsed.

We noted that reports of a prolapsed hemipenis are very rare, and our report is the first in a live, wild snake in North America, as well as the only report of this condition in *M. fuliginosus* (Podhade and Harne 2014). It is unknown what level of frequency this may be occurring in wild populations, but this phenomenon can be a confounding effect on the population persistence of declining species in some areas.

Acknowledgements—We thank to Maytorell family of the El Potrero Ranch for access to their site for investigations into the herpetofauna found there. We also acknowledge assistance in the field from: Diego Alvarado-Simon, Amanda Castro, Elias Elias, Andy Forde, Audrey Forde Andrew Isner, Heidi Isner, Robert Kely, Donna Noce, Ivan Parr, Anny Peralta-García, and Angel Ramirez. Collecting permit (SGPA/DGVS/01472/21) was issued by the Secretaría de Medio Ambiente y Recursos Naturales (SEMARNAT) to JHVV.

Literature Cited

Bennett, R.A., and D.R. Mader. 1996. Soft tissue surgery. Pages 287-298 *in*: D.R. Mader (editor). Reptile Medicine and Surgery. W.B. Saunders, Philadelphia, PA.

- Da Silva, K.B., D.F.M. Da Silva, K.M.P. Da Silva, V.C. Garcia, and S.M.A. Santos. 2013. Hemipenectomy in the golden lancehead (*Bothrops insularis*) – case report. *Veterinária e Zootecnia* 20:453-458.
- Hedley, J., and K. Eatwell. 2014. Cloacal prolapses in reptiles: a retrospective study of 56 cases. *Journal of Small Animal Practice* 55:265-268.
- Hernandez-Divers, S.J. 2004. Surgery: principles and techniques. Pages 147-167 *in*: P. Raiti and S. Girling (editors). *Manual of Reptiles*. British Small Animal Veterinary Association, Cheltenham, England.
- Kelly, D.A. 2002. The functional morphology of penile erection: tissue designs for increasing and maintaining stiffness. *Integrative and Comparative Biology* 42:216-221.
- Lescano, J., and M. Quevedo. 2016. Hemipenile prolapse and hemipenectomy in a leopard gecko (*Eublepharis macularius*). *Analecta Vet* 36:5-9.
- Nasoori, A., A. Taghipour, and B.A. Young. 2018. Male reproductive behaviour of *Naja oxiana* (Eichwald, 1831) in captivity, with a case of unilateral hemipenile prolapse. *Herpetology Notes* 11:1061-1064.
- Pernetta, A.P., and C.J. Reading. 2009. *Coronella austriaca* (smooth snake) Paraphimosis. *Herpetological Review* 40:95.
- Podhade, D.N., and R. Harne. 2014. Surgical intervention in prolapsed rescued Russell viper (*Daboia russelii*). *Journal of Wildlife Research* 2:9-10.
- Ortenburger, A.I. 1928. Whipsnakes and racers. Plimpton Press, Norwood, MA.
- Schultz, C., and A.J. Caven. 2021. Lined Snake (*Tropidoclonion lineatum*) prescribed fire mortality. *Transactions of the Nebraska Academy of Sciences and Affiliated Societies*. 530:42-45.
- Stebbins, R.C. 2003. Field guide to amphibians and reptiles of California. University of California Press, Berkeley, CA.
- Thomson, R.C., A.N. Wright, and H.B. Shaffer. 2016. California amphibian and reptile species of special concern. University of California Press, Berkeley, CA.
- Vyas, R., and V. Mistry. 2021. Male genital/cloacal prolapse in wild marsh crocodiles *Crocodylus palustris*, Gujarat, India. *The Herpetological Bulletin* 157:27-28.



We noted that reports of a prolapsed hemipenis are very rare, and our report is the first in a live, wild snake in North America, as well as the only report of this condition in *M. fuliginosus*. It is unknown what level of frequency this may be occurring in wild populations, but this phenomenon can be a confounding effect on the population persistence of declining species in some areas.